

LIGHTWAVE™ Therapy Consent Form

You have chosen to undergo LIGHTWAVE Therapy for the following purpose(s):

Items to consider before LIGHTWAVE™ Therapy:

- Review the other side of this document for contraindications. Notify your esthetician if any of the conditions or medications are applicable to you.
- Final results may not be apparent for several weeks after the treatments.
- Sun exposure, alcohol consumption, smoking, and eating habits directly affect treatment outcome. Remember to eat well, limit sun exposure, and keep alcohol consumption and smoking to a minimum.
- Drink at least 8 oz of water before and after each treatment.
- Do not apply creams, moisturizers, or antiperspirants to treatment area prior to the treatment.
- It is important to notify your esthetician if there are any problems or concerns after the treatment, including prolonged redness.
- The treatment fee is for provision of the treatment. There is no guarantee that every individual will achieve the expected or anticipated results. Individuals are, in fact, individuals after all.
- More than one treatment package may be required to achieve desired result.

My signature verifies that I have read and understand the goals, limitations, risks, and possible side effects to the treatment, and I have been given the opportunity to ask questions. My signature also verifies my informed decision to proceed with LIGHTWAVE™ Therapy and have the treatment.

Client Signature

Date

Parent/Legal Guardian (if Client under 16 years)

Date

I have fully explained to the Client, _____, the nature, purpose, and expected results of LIGHTWAVE™ Therapy and the risks involved. I have answered all questions regarding the treatment.

Esthetician

Date

Certain medications or medical conditions can cause a person to develop sensitivity to light. LIGHTWAVE has compiled a list of such medications and conditions; however, this is in no way intended to be a complete list of all possible conditions. If you are concerned about a medical condition or medication and its impact by LED, please consult with your physician prior to your first LIGHTWAVE™ Therapy treatment.

Acute or Cutaneous Porphyria	Migraines	Lupus Erythematosus
Heart Trouble/Pacemaker	Asthma	Thyroid Problems
Photophobia	Diabetes	Exogenous Eczema
Eye Diseases/Retinal Abnormalities	Pregnancy	Epilepsy or Seizures
Hypomelanism (albinism)	Skin Cancer	Blood Thinners

The following medications have been known to cause light sensitivity. It is recommended that the medications listed below be suspended for a minimum of one week before undergoing LIGHTWAVE™ Therapy. If it is not possible to discontinue the use of a medication, then you must consult your doctor before undergoing LIGHTWAVE™ Therapy while continuing the medication. Please also check with the prescribing physician before discontinuing any prescribed medications. This is a listing of the medications most commonly associated with photosensitivity; however, it is by no means a complete list of all photosensitive medications. Herbs and other over-the-counter medications can also cause sensitivity to light.

Acne: Oral Isotretinoin (Accutane, Accure, Aknenormin, Amnesteem, Ciscutan, Claravis, Isohexal, Isotroin, Oratane, Sotret, Roaccutane)
Topical Isotretinoin (Isotrex, Isotrexin)
Anti-Arrhythmic: Amiodarone (Pacerone, Cordarone, Aratac)
Chlorproazine (Thorazine, Chloramead, Chlordryprom, Chlor, Promanyl, Largactil, Promapar, Promosol, Terpium, Sonazine)
Antibiotics: Tetracycline (Helidac, Terra-Cortril, Terramycin, Sumycin, Actisite, Bristacycline, Actisite, Tetrex, Doxycycline, Ciprofloxacin)
Norfloxacin (Noroxin, Quinabic, Janacin)
Ofloxacin (Floxin, Oxaldin, Tarivid)
Nalidixic Acid (NegGam, Wintomylon)
Ciprofloxacin (Cipro, Ciproxin, Ciprobay)
Minocycline (Minomycin, Minocin, Arestin, Akamin, Aknemin,)Solodyn, Dynacin, Sebomin)
Oxytetracycline, Demeclocycline, Lymecycline
Antifungal: Griseofulvin (Grifulvin)
Antipsychotic: Haloperidol (Haldol)
Trifluoperazine (Stelazine, Clnazine, Novoflurazine, Pentazine, Solazine, Terfluzine, Triflurin, Tripazine)
Cancer: Methotrexate (MTX, Aminopterin, Ledertrexate)
Arthritis: Auranofin (Ridaura) If taking this medication, you are not a candidate for light therapy.